

SUMAROKOV, V.P.; VOLODUTSKAYA, Z.M.

Complex esterification of tall oil acids with isoamyl alcohol.
Gidroliz. i lesokhim. prom. 17 no.3:11-13 '64.

(MIRA 17:9)

1. NITKhIB.

SUMAROKOV, V.P.; VOIODUTSKAYA, Z.M.

Production of high-quality methylacetate. Gidroliz. i lesokhim.
prom. 17 no.7:16-17 '64. (MIRA 17:11)

1. NITKhIB.

SUMAROKOV, Viktor Pavlovich VAN'YAN, Mariya L'vovna; ASKINAZI,
Anna Il'inichna; TULYAKOV, B.V., red.

[Tall oil] Tallovoe maslo. Moskva, Lesnaia promyshlennost',
1965. 146 p. (MIRA 18:3)

SUMARCKOV, V.P.

Сумарков, В.П.

New standard for wood alcohol solvents. Gidroliz. i lesokhim.
prom. 18 no.3:6-7 '65.

(MIRA 18:5)

KONDRAT'YEVA, Ye.N.; TARANENKO, I.I.; SUMARUKOVA, R.S.

Requirement of some microelements by purple and green sulfur
bacteria. Nauch. dokl. vys. shkoly; biol. nauki no.2:176-180
'65. (MIRA 18:5)

1. Rekomendovana kafedroy mikrobiologii Moskovskogo gosudarstvennogo
universiteta im. M.V. Lomonosova.

LITVYAK, I.G.; SUMAROKOVA, T.N.

Solvolysis of tin tetrahalides. Zhur. ob. khim. 34 no.11:
3677-3682 N '64 (MIRA 18:1)

SUMAROKOV, V. S.

Hydrology

DECEASED

C. 162

1767

191T18

SUMAROKOVA, M.

USSR/Chemistry - Antimony Compounds Jul 51

"Binary Systems Formed by SbCl_3 , SbCl_5 , and AsCl_3
V. The System SbCl_3 - CCl_3COOH ," T. Sumarokova,
M. Usanovich, Lab Phys Chem, Inst Chem, Acad Sci
Kazakh SSR

"Zhur Obshch Khim" Vol XXI, No 7, 1219-1222

Studied elec cond, viscosity, density of system
 SbCl_3 - CCl_3COOH at 50, 60, 70°C. Elec cond and
viscosity diagrams showed presence of acid-base
interaction between components and of compds
 SbCl_3 · CCl_3COOH and 2SbCl_3 · CCl_3COOH .

191T18

USSR/Chemistry - Antimony Compounds Jul 51
(Contd)

Dyatectic max on fusibility diagram of system
represented compd SbCl_3 · CCl_3COOH (mp 56°).
This is 1st established case of CCl_3COOH acting
as additive or oxonium base.

191T18

SUMAROKOVA, M.Ya., dotsent.

Work of the scientific student circle attached to the chair of pharmacognosy
of the Chervenkov Academy of Medicine in the academic year 1951-52 (Bulgaria).
Apt.delo 2 no.5:51-52 S-0 '53. (MLRA 6:10)
(Bulgaria--Pharmacognosy--Study and teaching)
(Study and teaching--Pharmacognosy--Bulgaria)

SUMAROKOVA, M.Ya., dotsent; DAVRILYUK, Ye.Ya.

From the pages of the foreign pharmaceutical press (Czechoslovakia). Apt.
delo no.4:74-79 J1-Ag '53. (MLBA 6:8)

(Czechoslovakia--Pharmacy) (Pharmacy--Bibliography)

(Bibliography--Pharmacy)

SUMAROKOVA, M.Ya., dotsent

Fifth edition of the Hungarian Pharmacopeia, 1954. Apt. delo 3
no.5:54-57 S-O '54. (MIRA 7:12)
(PHARMACOPEDIA,
Hungar, 5th edition)

SUMAROKOVA, M.Ya., dotsent.

Pharmaceutical education in Czechoslovakia. Apt.delo 4
no.5:59-64 S-O '55. (MLBA 8:12)
(PHARMACY, education,
in Czech.)

SUMAROKOVA, M.Ya., dotsent.

First supplement to the seventh edition of the French Pharmacopoeia.
M.Ia. Sumarokova. Apt. delo. 4 no.6:50-52 N-D '55. (MIRA 9:1)

(FRANCE-PHARMACOPOEIAS)

SUMAROKOVA, M.Ya.,dotsent

"Pharmacy." Reviewed by M.IA.Sumarokova. Apt.delo 5 no.2:57-59
Mr-Ap 256. (MLRA 9:7)
(CZECHOSLOVAKIA--PHARMACY--PERIODICALS)

GOLOSOV, A.V.; SOKOLOV, I.I.; USPENSKAYA, A.N.; TSVETKOV, N.G.; SUMAROKOVA,
M.Ya., redaktor; CHERNYAVSKIY, M.N., redaktor; LYUDKOVSKAYA, N.I.,
technicheskiy redaktor.

[Textbook of the Latin language for secondary medical schools]
Uchebnik latinskogo iazyka dlia srednikh meditsinskikh uchebnykh
zavedenii. Pod obshchei red. M.IA.Sumarokovoi. Moskva, Gos.isd-vo
med.lit-ry, 1957. 157 p. (MIRA 10:11)
(Latin language)

SUMAROKOVA, M.Ya., dotsent

Pharmaceutical education in various countries. Apt.delo 6 no.1:
80-83 Ja-Y '57. (MLRA 10:3)
(PHARMACY--STUDY AND TEACHING)

SUMAROKOVA, M.Ya., dotsent

The United States Pharmacopoeia. Fifteenth Revision, 1955;
brief survey. Apt.delo 6 no.2:87-89 Mr-Ap '57. (MLRA 10:6)
(UNITED STATES--PHARMACOPOEIAS)

Sumarckova, M.Ya.
SUMARCKOVA, M.Ya., dots.

~~Finnish pharmacopoeias, seventh edition, 1956. Apt.delo 6 no.3:~~
77-80 My-Je '57. (MIRA 11:1)
(FINLAND--PHARMACOPOEIAS)

SUMAROKOVA. M.Ya., detsent

"For health protection" ("Au service de votre santé." Reviewed by
M.I.S. Sumarokova). Apt. 6 no. 69-72 Ul-Pr '57. (MIRA 10:9)
(FRANCE--PHARMACY)

SUMAROKOVA, M.Ya., dotsent

Material on the training of pharmacists in a number of countries.

Apt.delo 6 no.6:78-80 N-D '57.

(MIRA 10:12)

(PHARMACY--STUDY AND TEACHING)

SUMAROKOVA, M.Ya., dotsent

Some problems of pharmacy in the United States; from material
published in the Journal of the American Pharmaceutical Association,
February 1958. Apt.delo no.4:83-85 J1-Ag '58 (MIRA 11:8)
(UNITED STATES--PHARMACY)

SUMAROKOVA, M.Ya., dots.

Some information on the preparation of the new DAB-7 edition of
the German pharmacopoeia. Apt.delo 7 no.5:88-89 3-0 '58 (MIRA 11:10)
(GERMANY---PHARMACOPOLIAS)

SUMAROKOVA, M. Ya., dots.


Pharmacopoeia of the Netherlands for 1958. Apt.delo 8
no.2:88-89 Mr-Ap '59. (MIRA 12:5)
(NETHERLANDS--PHARMACOPOEIAS)

AUTHORS: Kolchin, O.P., Sumarokova, N. V., Chuveleva, N. P., 89-12-5/29


TITLE: Production of Plastic Niobium (Poluchenije plastichnogo niobiya)

PERIODICAL: Atomnaya Energiya, 1957, Vol. 3, Nr 12, pp. 515-524 (USSR)

ABSTRACT: First the properties of niobium are written down. The process necessary for obtaining niobic powder with 98,9 - 99,2% content of niobium is described in detail. This powder is obtained by reduction of the K_2NbF_7 with sodium. If from the powder obtained pressed bars are manufactured and sintered in the vacuum, plastic niobium is obtained. The investigation of the phase condition of this not entirely reduced mixture of niobic oxide and niobic carbide shows that the main reaction in the reduction in the vacuum can be understood as a summation reaction of successively occurring reactions in the following form:



where "NbC" is the phase with variable composition:



Special investigations gave evidence that in the choice of the reduction regimen the interaction between the vapours of the inter-

Card 1/2

S/180/61/000/006/002/020
E021/E135

AUTHORS: Sazhin, N.P., Kolchin, O.P., and Sumarokova, N.V.
(Moscow)

TITLE: The processes of reduction of niobium oxides by
carbon

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Otdeleniye
tekhnicheskikh nauk. Metallurgiya i toplivo,
no.6, 1961, 8-24

TEXT: The chemical and physical processes occurring during
high temperature reduction of oxides of niobium by carbon were
studied with the aim of explaining the mechanism of reduction.
Niobium pentoxide powder (0.15 mm particles) containing 0.01% TiO_2 ,
0.06% Fe_2O_3 and 0.01% SiO_2 was used with lamp-black or niobium-
carbide as reducing agents. The niobium carbide was prepared by
heating a mixture of niobium carbide with lamp black at 1800 °C in
a current of hydrogen; it contained 10.2% carbon. The phase
composition of the products of incomplete reduction were studied,
a thermodynamic analysis of the Nb-O-C system was made and the
properties of the oxides of niobium were investigated. The rates
Card 1/4

The processes of reduction of ...

S/180/61/000/006/002/020
EO21/E135


of diffusion of oxygen and carbon in niobium and the rates of evaporation of the lower oxides of niobium were compared semi-quantitatively. From the results of the experiments and from a critical examination of other literature it is shown that the reduction is a multi-stage process, and a mechanism for reduction at temperatures used in practice is proposed. Reduction by niobium carbide at 1100-1300 °C of the pentoxide to the dioxide and partially to the oxide occurs in the main by the generally accepted two-stage scheme with the formation of carbon monoxide. Reduction may also occur at the beginning of the process by interaction in the solid phase between niobium pentoxide and carbon formed by dissociation of the carbide. The second stage of the process is the reduction of the lower oxides of niobium at temperatures above 1500 °C. This takes place in the gaseous phase and depends on the evaporation of the lower oxides which is the reaction limiting the rate of the process. In the last stage of the process, oxygen and carbon diffuse to the surface of the pores in the metal and are evolved as carbon monoxide. The reaction limiting the purification of the metal from dissolved

Card 2/4

The processes of reduction of ...

S/180/61/000/006/002/020
E021/E135

oxygen and carbon is the desorption of the carbon monoxide. The reduction of the pentoxide to the dioxide and the dioxide to the oxide can be carried out in a relatively low vacuum or in a current of hydrogen or inert gas. The reduction of the oxide and the removal of the carbon monoxide at temperatures used in practice require a high vacuum. The rate of reduction can be increased by increasing the temperature and rate of carbon monoxide removal. The maximum temperature possible for each stage is determined by the melting point of the phase most easily melted. The proposed mechanism for the reduction process of niobium can be extended without any radical changes to the reduction process of tantalum and vanadium from their oxides and from mixtures of their oxides and carbides. There are 6 figures, 4 tables and 33 references: 19 Soviet-bloc, 6 Russian translations from non-Soviet publications, and 8 non-Soviet-bloc. The four English language references read as follows:



Card 3/4

The processes of reduction of ...

S/180/61/000/006/002/020
E021/E135

Ref. 15: G. L. Miller. Tantalum and Niobium, London, 1959,
pp 181-187, 283-291.

Ref. 17: High Temperature Technology. N.Y.-London, 1956,
ed. J. E. Campbell.

Ref. 24: F. Holtsberg, A. Reisman, M. Bewry, M. Berkonbilt.
The polymorphism of Nb_2O_5 . J. Amer. Chem. Soc., 1957,
79, 2039.

Ref. 26: R. Orr. High temperature heat contents of tantalum and
niobium oxides. J. Amer. Chem. Soc., 1955, 75, 2808-09.

SUBMITTED: March 21, 1961

Card 4/4

S/089/61/010/002/012/018
B102/B209

AUTHORS: Kolchin, O. P., Sumarokova, N. V.

TITLE: Melting point and other properties of lower niobium oxides

PERIODICAL: Atomnaya energiya, v. 10, no. 2, 1961, 168-170

TEXT: The present paper is a report on investigations made on NbO and NbO₂. NbO₂ was produced by reduction of the pentoxide in vacuo at 1300 and 1700°C, NbO by reduction of NbO₂. The impurities in the pentoxide (tantalum-silicon-, titanium-, and iron oxides) were removed for the major part during reduction in vacuo. The three obtained niobium dioxide samples still contained 0.02 - 0.04% by weight of C, 0.006% by weight of N and had the composition NbO_{1.942}, NbO_{1.956} and NbO_{1.986}. Radiographic structural analysis showed that the product consisted of one phase only with the following lattice parameters: $a = 4.82 \pm 0.02$ kX, $c = 2.99 \pm 0.02$ kX. The three monoxide samples contained 0.04 - 0.06% by weight C and 0.03% by weight N and exhibited the composition NbO_{0.95}, NbO_{1.01}, and NbO_{1.02}. Melting point determination: Briquets

Card 1/3

Melting point and other ...

S/089/61/010/002/012/018
B102/B209

(1.5 - 2.0 t/cm², 10 x 10 mm cross section) were pressed from NbO and NbO₂ powder and electrically heated in containers made of niobium sheet by a graphite heater in vacuo. The melting point of NbO was found to be at 1935°C, that of NbO₂ at 2080°C, allowing for an error of $\pm 15^\circ\text{C}$. Micro- and radiographic structural analysis was employed on re-molten NbO for checking the single-phase consistency and to determine the parameter a (4.201 kX). The results concerning NbO₂ do not agree with those from Ref. 9 which is said to be due to the fact that in that case NbO₂ contained NbO impurities. Determination of volatility and composition of the gaseous phase: Investigations in vacuo ($0.5 - 1 \cdot 10^{-6}$ mm Hg) at various temperature showed that both oxides start evaporating at a considerable rate at 1700°C. At 1850°C, all of the monoxide and 45% by weight of the dioxide were evaporated after 4 hours, and after 8 hours also the dioxide was entirely evaporated. It was proved mass-spectroscopically that the gaseous phase above NbO₂ contains NbO₂ molecules only, i. e. that neither dissociation nor association take place. Investigation of the micro-hardness: Re-molten homogeneous niobium oxide samples had a micro-hardness of

Card 2/3

Melting point and other ...

S/089/61/010/002/012/019
B102/B209

1930 kg/mm², the eutectic in the average one of 794 kg/mm² (load of 50 g), the dioxide 1720 kg/mm²; the latter value is inaccurate. Character of electric conductivity: NbO turned out to have metallic conductivity, NbO₂ to be a semiconductor. Oxidation in air: Dioxides (NbO_{2.00} and NbO_{1.92})² produced at 1300 and 1700°C, respectively, and a monoxide (NbO_{0.914}) produced at 1700°C were powdered (grain size 0.15mm) and heated in air at 100 - 300°C; oxygen content was determined. After 6 hours of heating at 100, 150, 200, 225, 250, and 275°C the oxygen content was unchanged in the monoxide and was a little raised in the dioxide at 275°C. However, the surface of the powder particles of both oxides became yellowish already at 150°C and bronze-colored at 200°C. Only after 6 hours at 300°C, both oxides were completely oxidized to the pentoxide. In conclusion, the authors thank L. V. Mel'nikova for having made the metallographic analysis. There are 2 figures and 11 references: 5 Soviet-bloc and 2 non-Soviet-bloc. ✓

SUBMITTED: October 4, 1960

Card 3/3

ACCESSION NR: AP4042350

were sintered at a temperature varying from 1700 to 2100C; compacts 20 x 20 mm were sintered at 1900C or at 2100C in a vacuum of 0.001 mm Hg. In the preparation of alloys, Al, Ti, Zr, V, Ta, Mo, and W were used as the alloying elements; for more complete removal of carbon during reduction, the oxides were added in an amount 2—5% higher than the stoichiometric. The experiments showed that binary Nb—(4.8—6.2)%Mo, Nb—(5.7—24.6)%W, Nb—(4.3—5.8)%V, Nb—2.2% Ti alloys, ternary Nb—(4.25—4.9)%Mo—(0.87—1.85)%Zr, Nb—(17.0—26.4)%W—(2.07—4.5)%Ti, Nb—5.0%W—2.0%Ta, Nb—(3.0—3.5)%V—(0.4—0.5)%Al alloys, and quaternary Nb—14.1%W—5.0%Mo—(0.93—1.1)%Zr alloys can be prepared by one or both processes. Attempts to obtain binary Nb—Al alloys were unsuccessful. Better quality alloys with a consistently lower content of O, N, and C are obtained by reduction at 2100C. The alloys have a porosity of 40—50%. Subsequent electron beam melting substantially lowers the content of O, N, and C, in some cases without affecting the content of W and Mo, or Zr when its content is about 1%. When necessary, the alloys can be reprocessed by any conventional method used for unalloyed niobium. Experiments on electron beam melting of the alloys were conducted by A. V. Yelyutin. Orig. art. has: 3 tables.

Card 2/3

ACCESSION NR: AP4042350

ASSOCIATION: none

SUBMITTED: 00

SUB CODE: MM

ATD PRESS: 3069

ENCL: 00

NO REF SOV: 004

OTHER: 002

Card 3/3

L 32686-66 EWT(m)/T/EWP(t)/ETI IJP(c) JD/JG

ACC NR: AP6012727

SOURCE CODE: UR/0136/66/000/004/0067/0070

AUTHOR: Kolchin, O. P.; Sumarokova, N. V.; Vol'dman, M. A.

ORG: none

TITLE: Kinetics of the combined carbothermic reduction of niobium and tungsten

SOURCE: Tsvetnyye metally, no 4, 1966, pp 67-70

TOPIC TAGS: vacuum furnace, chemical reduction, niobium, tungsten, niobium compound/
/VVPS-10A type vacuum furnace

ABSTRACT: This is a continuation of a previous investigation (O. P. Kolchin et al. Tsvetnyye metally, 1964, no 7) with the difference that it deals with a detailed investigation of the kinetics of the combined carbothermic reduction of Nb and W from the mixtures of the oxides and carbides of Nb and alloy elements at various temperatures, the degree of reduction being determined according to the change in the C content of specimens following their heating in a VVPS-10A type vacuum furnace. In the reduction products the W content was determined by the photolorimetric thiocyanate method, correct to 3-5% (rel.); the Nb content, according to weight gain when heating the specimen in air; and the C content, by the volumetric method. It was found that in the presence of W the degree of the reduction of Nb_2O_5+5NbC at 1200 and

Cord 1/2

UDC: 669.293'27.094.2

L 32686-66

ACC NR: AP6012727

1400°C is greater, particularly so when the raw charge contains 30% W in the form of an oxide rather than a carbide. X-ray structural phase analysis showed that W is the first to get reduced and subsequently, at higher temperatures, it forms an alloy with Nb, which is accompanied by the reduction of Nb. It is best to start with the temperature of 1400°C and regularly increase it until at most 1900°C. These findings confirm the earlier observation that W oxides during the reduction virtually do not get sublimated from the burden and that even alloys with a high W content have a close-to-theoretical composition. Orig. art. has: 1 figure, 1 table,

SUB CODE: 11, 13

SUBM DATE: none/ ORIG REF: 004

Card 2/2 BLG

L 40919-66		ENP(e)/ENT(m)/ENP(l)/ETI/ENP(k)		ISF(c)	JD/JJ
ACC NR: AP6020738		SOURCE CODE: UR/0136/66/000/006/0065/0067			
AUTHOR: <u>Kolchin, O. P.</u> ; <u>Chuveleva, N. P.</u> ; <u>Sumarokova, N. V.</u> ; <u>Filipenko, V. V.</u> ; <u>Men'shchikov, V. A.</u> ; <u>Kadyshevskiy, V. S.</u> ; <u>Belimov, N. I.</u> ; <u>Abramovich, E. B.</u>					
ORG: none					
TITLE: Manufacture of powdered niobium and its alloys by hydrogenating compacted metals and alloys					
SOURCE: Tsvetnyye metally, no. 6, 1966, 65-67					
TOPIC TAGS: metal powder, powder metal production, niobium, powder metallurgy, hydrogenation, niobium alloy					
ABSTRACT: The report presents a method for manufacturing high purity powders by hydrogenating niobium or its alloys at lower temperatures (360 to 400C) and lesser excess hydrogen pressures (up to 0.7 atm) than those commonly required. The process is even faster at the reduced temperature levels. Hydrogenation and milling techniques are given in detail for source materials derived by electron beam smelting or carbide heating processes. For the latter, direct yield of dehydrogenated powder was 91.4%, total yield 93.3%, unaccountable losses 1.1%. The impurity content in niobium powders obtained from different compacted metals is					
Card 1/3		UDC: 669.293-492.2			

L 40919-66

ACC NR: AP6020738

given in Table 1.

Table 1. Impurity content (% by mass) in niobium powders obtained from different compacted metals.

Initial material			Powder (-0.147 mm)		
N	O	C	N	O	C
Reduced Metal					
0.04	0.27	0.15	0.04	0.24	—
0.05	0.27	0.03	0.05	—	—
0.05	0.20	0.09	0.05	—	0.11
0.05	0.20	0.10	0.03	—	—
0.04	0.23	0.07	0.06	—	—
0.04	0.13	0.06	0.09	—	—
0.07	0.24	0.05	0.05	0.32	—
0.05	0.20	0.07	0.04	0.30	—
0.05	0.15	0.06	0.05	—	—
Ends of rods of a sintered Metal*					
0.05	—	0.12	0.08	0.16	0.15
0.04	0.45	0.20	0.02	0.46	0.25
0.05	0.25	0.12	0.05	—	0.11
0.04	0.27	0.03	0.05	0.30	0.11
—	—	—	0.05	0.35	0.36
0.05	—	—	0.06	0.40	0.20

*The sintered rods contain 0.01-0.03% C; 0.02% N; 0.02% O; <0.01% Ti; 0.01-0.03% Si; 0.01-0.03% Fe; 0.15-0.25% Ta; ~59.5% Nb (+Ta).

Card 2/3

L 40919-66

ACC NR: AP6020738

Orig. art. has: 2 figures and 1 table.

SUB CODE: 11,13/ SUBM DATE: 00/ ORIG REF: 001/ OTH REF: 002

Card 3/3 11b

L 0007-67 EST(D)/T/EMP(C)/RTI LIP(C) JD/JG

ACC NR: AP6031728

SOURCE CODE: UR/0136/66/000/009/0072/0074

AUTHOR: Kolchin, O. P.; Filipenko, V. V.; Nizharadze, K. S.; Abramovich, E. B.; Sumarokova, N. V.; Men'shchikov, V. A.

ORG: none

TITLE: Synthesis of niobium carbide with a low nitrogen content

SOURCE: Tsvetnyye metally, no. 9, 1966, 72-74

TOPIC TAGS: niobium carbide, high purity carbide, ~~nitrogen niobium carbide~~, niobium carbide synthesis, niobium compound, carbide, nitrogen, oxygen, chemical synthesis

ABSTRACT: An investigation has been made of the various factors which contribute to the contamination with nitrogen and oxygen of niobium carbide produced by a continuous process in the Tamman furnace. The investigation results showed that the only significant source of contamination was the inflow of air into the reaction chamber when the furnace was opened every 30 min for charging and removing the final product. Modification of the charge chamber decreased the cross section of the charging shute from 1000 to 160 cm², cut in three the number of openings required to charge the chamber, and sharply reduced the amount of the air flowing in through a narrowed charge shute. A hydraulic lock was also installed for combustion gases, which made it possible to increase the pressure of gases in the furnace to 100—200 mm Hg and thus practically eliminate the inflow of air into the furnace.

Card 1/2

UDC: 669.293

L 0007-67

ACC NR: AP6031728

APPROVED FOR RELEASE: 08/26/2000 **CIA-RDP86-00513R001653910020-4**
The resulting improvement in the quality of niobium carbide produced. The niobium carbide produced in the modernized furnace contained 89.32—89.63% Nb(+Ta), 0.03—0.14% Fe, 10.0—10.4% C, and only 0.028—0.059% N and 0.14—0.52% O, instead of the previous 0.3% N and 2—3% O. Tantalum carbide with a low content of nitrogen and oxygen was also produced in the modernized furnace, and it is believed that pure carbides of other refractory metals can be produced in it. Orig. art. has: 2 figures.

SUB CODE: 07 / SUBM DATE: none / ORIG REF: 005 / OTH REF: 001

Card

2/2

137 AND 138 CODES) PROCESSES AND PROPERTIES INDEX

AC 7-1

Conductivity, viscosity, and surface tension in the system $H_2O-HClO_4$. M. URANOVITSCH, T. SUMAROKOVA, and V. UDOVERKO (J. Gen. Chem. Russ., 1939, 8, 1167—1975).—Conductivity, η , and surface tension data for the system at 0°, 10°, and 25° do not suggest compound formation. The max. on the conductivity-composition curve is ascribed to lowering of the η of the system with increasing $[HClO_4]$. R. T.

ASA-SLA METALLURGICAL LITERATURE CLASSIFICATION

CA

The electric conductivity, viscosity and surface tension of the system $H_2SO_4-HClO_4$. M. Usmorovich, T. Sumarokova and V. Udevnenko. *Acta Physicochem. U.R.S.S.* 11, 505-509(1958) (in German); *J. Gen. Chem. (U.S.S.R.)* 28, 201-202(1958).—An app. for the simultaneous detn. of the elec. cond., viscosity and surface tension is described. Pure $HClO_4$ was prepd. by distn. of 70% acid with 98% H_2SO_4 . The phys. consts. were detd. at 0°, 10° and 20°, resp.: $\kappa \times 10^3$, 3.619, 3.385, 4.063;

0, 10° and 30°, resp.: $\chi \times 10^4$, 11.20, 9.68, 7.95; d , 1.2129, 1.7918, 1.7722. Data are given on the χ , ν , ϕ , and d values of various $\text{HClO}_4\text{-H}_2\text{SO}_4$ azeot. The ν ($\text{mol. } \% \text{H}_2\text{SO}_4$) curves rise slowly up to 30 mol. % H_2SO_4 , then rapidly to 100% H_2SO_4 : $\chi \times 10^4$ for 37.5% H_2SO_4 = 4.42, 3.84 and 2.68 at 0°, 10° and 25°, resp.; for 50.75% H_2SO_4 , 10.20, 7.60, 4.31. The $\nu = f(\text{mol. } \% \text{H}_2\text{SO}_4)$ curves pass through a max. at about 33% H_2SO_4 , $\chi \times 10^4$ = 14.5, 18.5, 18.6. The $\chi\phi$ curves are convex to the mol. % axis. The ϕ curves are almost linear but slightly concave, and obey fairly well the Whittleworth equation $\phi = [c_1X + (1-X)c_2]/R$, where X is the mol. fraction of H_2SO_4 and R is the correction factor calcd. from the d . curve (R is a min. at 30 mol. % = 0.9604). It is concluded that the 2 components of the system do not interact with one another.

F. H. Rathmann

F. H. Rathmann

ALB-114 METALLURGICAL LITERATURE CLASSIFICATION

Electric conductivity, viscosity, and density of binary systems with perchloric acid. II. The system $\text{HClO}_4\text{-CCl}_3\text{COOH}$. *I. Sumerukhov and Z. Grushkin. J. Gen. Chem. (U.S.S.R.)* 16, 1991-6 (1946) (in Russian); *J. Gen. Univovich, C.A.* 34, 31567.---Absence of acid-base interaction and behavior of CCl_3COOH as indifferent solvent towards HClO_4 in complete analogy with its behavior towards H_2SO_4 (cf. the foregoing abstr.), follows from facts of κ , η , and d , at 20, 30, and 60°. Viscosity η rises continuously from HClO_4 to CCl_3COOH ; the curves are convex to the axis of compn.; the decrease of η at higher temp. is more pronounced at higher CCl_3COOH contents; curves of κ have a similar shape but fall uniformly and rapidly from HClO_4 to CCl_3COOH ; the temp. coeff. of the elec. cond. (between 30 and 60°) rises rapidly from HClO_4 to CCl_3COOH ; example of data, at 60°, HClO_4 79.7, 34.8, 23.7 mol. %, η = 0.00793, 0.01811, 0.02366, κ = 0.00366, 0.00128, 0.00086, ρ = 0.79, 4.63, 11.45%. The κ curve falls almost linearly (with a slight convexity to the axis of compn.) from HClO_4 to CCl_3COOH . The sp. vol. V_d rises from HClO_4 to CCl_3COOH along a curve concave to the axis of compn.; example, at 60°, HClO_4 79.7, 34.4, 23.7 mol. %, d = 1.6360, 1.6335, 1.6177.

N. Thon

N. Thon

A.S.M.-I.I.A. METALLURGICAL LITERATURE CLASSIFICATION

SUMAROKOVA, T.

USSR/Electricity
Conductivity
Conductance - Charts

Sep/Oct 46

"The Electrical Conductivity of the System $\text{HClO}_4 \cdot \text{H}_2\text{O}$," M. Usanovich, T. Sumarokova,
Lab Phys Chem, Middle Asia State U, Tashkent, 5 pp

"Axta Physicochimica URSS" Vol XXI, No 5

Complete electrical conductance diagrams obtained at temperatures 20° , 50° , 60° ; and the 50° isotherm investigated at great length. Data obtained indicates chemical interaction in system; hydrates of perchloric acid, $\text{HClO}_4 \cdot \text{H}_2\text{O}$ appear to manifest themselves in liquid phase. Received, 1 Aug 1945.

PA 54T45

SUMAROBova, T.

PA 54T38

USSR/Chemistry - Acetic Acid
Chemistry - Perchloric Acid

Sep/Oct 1946

"The Systems Formed by Perchloric Acid With Acetic Acid and Its Chlorine Derivatives," T. Sumarobova, M. Ulanovich, Lab Phys Chem, Middle Asia State U, Tashkent, 8 pp

"Acta Physicochimica URSS" Vol XII, No 5

Experiment undertaken to display possible basic properties in CCl_3COOH . Electrical conductivity, viscosity and density of systems HClO_4 - CHCl_2COOH , HClO_4 - CCl_3COOH , HCCl_3 - ClCOOH , HClO_4 - CH_3COOH were studied, and it was established that HClO_4 does not form any complex compounds with CCl_3COOH . The compounds

~~1946~~

54T38

USSR/Chemistry - Acetic Acid (Contd) Sep/Oct 1946

CHCl_2COOH , 2HClO_4 , CHCl_2COOH , HClO_4 , CH_2ClCOOH , $2\text{CH}_3\text{COOH}$, HClO_4 were detected; formation of these compounds is indicative of the basic properties of CHCl_2COOH , CH_2ClCOOH , CH_3COOH . Received, 1 Aug 1945.

54T38

14

Electroconductivity, viscosity, and density of binary systems formed by HClO_4 . III. The system $\text{HClO}_4\text{-CHCl}_3\text{COOH}$. T. Samarkova and M. Usanovich. *J. Gen. Chem.* (U.S.S.R.) 17, 157 (2/1947) (in Russian); cf. *C.A.* 41, 6796g, 7216a. Viscosity-concn. curves are S-shaped, values for HClO_4 and for CHCl_3COOH at 20° being 0.00481 and 0.0129, at 35°, 0.00889 and 0.0260; and at 50°, 0.01312 and 0.01790 poises. The cond.-concn. curves rise to rather sharp max. at a concn. of HClO_4 of 79.5 mole %, the values of the max. cond. at 20, 35, and 50° being 0.02587, 0.02629, and 0.02618 $\text{ohm}^{-1}\text{cm}^{-1}$. The curve of ρ d. against concn. is slightly S-shaped, deviating somewhat from a straight line. Values of ρ d. for HClO_4 and CHCl_3COOH at 20° are 1.778 and 1.526 g./cc. The data are interpreted to indicate the existence of two binary compds. with the formulas $2\text{HClO}_4\text{-CHCl}_3\text{COOH}$ and $\text{HClO}_4\text{-CHCl}_3\text{COOH}$, with CHCl_3COOH being basic

2

relative to the HClO_4 . IV. The system $\text{HClO}_4\text{-CH}_3\text{-CICOOH}$. M. Usanovich and T. Samarkova. *Ibid.* 163-4. Viscosity-concn. curves show max. at a concn. of HClO_4 of about 32.7 mole % at 35 and 50°, but crystals of CH_3CICOOH prevents observation of max. at 20°. Values of max. viscosity at 35 and 50° are 0.05981 and 0.02441 poises. Cond. curves show max. at about 80 mole % HClO_4 , values for cond. of HClO_4 being 0.02250, and of CH_3CICOOH less than 0.001, 'very small temp. dependence), and the max. cond. at 20, 35, and 50° being 0.02261, 0.02300, and 0.02778 $\text{ohm}^{-1}\text{cm}^{-1}$. The curve of temp. coeff. against HClO_4 concn. is S-shaped, rising from practically 1.0 for 100 mole % HClO_4 to about 4.0 below 25 mole % HClO_4 . The curve of ρ d. against concn. is convex towards the concn. axis; at 20°, e.g., it rises from 0.87 for 100 wt. % HClO_4 to 0.66 for 20 wt. % HClO_4 . The formation of one compd. is indicated, having the formula $\text{HClO}_4\text{-CH}_3\text{CICOOH}$, the latter being basic relative to the former. Arild J. Müller

СУМАРОВА, Т.

Usanovich, M., and Sumarokova, T. - "Electroconductivity, Viscosity and Density of Binary Systems of Binary Systems formed by HClO_4 . IV. The System HClO_4 — CH_2ClCOOH ." (p. 168)

SO: Journal of General Chemistry, (Zhurnal Obshchei Khimii), 1947, Vol. 17. No. 2.

Electroconductance, viscosity, and density of the binary systems formed by perchloric acid. V. System perchloric acid-acetic acid. M. Usanovich and T. Sumarokova. *Zhar. (Mokh. Khim. (J. Gen. Chem.)* 17, 1115-21 (1947); cf. C.A. 42, 3651f. -- Measurements were made at 20, 35, and 50°. The viscosity-concn. curves at all temps. studied have max. at 32.54 mole % HClO_4 . The electrocond. concn. curves at all 3 temps. rise sharply from pure HClO_4 to a max. at 82.85 mole % HClO_4 , then drop to a min. at 32.34%, rise to a slight max. at 12.11%, and then drop to the low cond. of pure AcOH . Values of d decrease steadily from pure HClO_4 to pure AcOH , values at 20°, e.g., being: HClO_4 , 1.7716, 51.17 mole % HClO_4 , 1.5746; 4.66 mole % HClO_4 , 1.1176. Data indicate the formation of the compd. $\text{HClO}_4 \cdot 2\text{AcOH}$, with no indication of the previously investigated compd. $\text{HClO}_4 \cdot \text{AcOH}$. VI. System perchloric acid-water. *Ibid.* 1122-7. -- Measurements were made at 20, 50, and 60°. The cond.-concn. curves rise from the cond. of pure HClO_4 to a slight max. at 54.3 mole % HClO_4 , then rise rapidly to a sharp max. at 10.10 mole % HClO_4 , and then drop sharply to the cond. of pure water. The viscosity-concn. curves go through a max. at 6.87 (centipoises) at 42.8 mole % HClO_4 . No d data are given in the article. The results indicate the formation of the compds. $\text{HClO}_4 \cdot \text{H}_2\text{O}$ and $\text{HClO}_4 \cdot 2\text{H}_2\text{O}$.
Arlid T. Miller

SUMAROKOVA, T.

Usanovich, M., and Sumarokova, T.- "Electroconductivity, Viscosity and Density of the Binary Systems formed with FCIO_4 . VI. The System $\text{HClO}_4\text{—H}_2\text{O}$ (p. 1427)

SO: Journal of General Chemistry, (Zhurnal Obshchei Khimii), 1947, Vol. 17, No. 8

SUMAROKOVA, T.N.

Binary systems containing perchloric acid. Izv. AN Kazakh. SSR Ser.
khim. no.3:33-39 '49. (MLBA 9:8)
(Acids, Organic) (Perchloric acid)

Синтетические, минеральные.

Acids, Organic

Complex compounds SnCl_4 , SbCl_3 , and AsCl_3 with some organic acids, Izv. Sek. plat. i blag. met. No. 25, 1950.

9. Monthly List of Russian Accessions, Library of Congress, April 195²₂, Uncl.

SHARONOVA, T.

"Binary systems containing SnCl_4 , SbCl_3 , and AsCl_3 . I. The system $\text{SnCl}_4\text{-CH}_2\text{ClCOOH}$." by N. Usanovich, T. Sharonova, and V. Glushchenko. (p.981)

SO: Journal of General Chemistry (Zhurnal Obshchei Khimii) 1951, Voluem 21, No. 6

SHIMIZU, T.

"Binary systems containing SnCl_4 , SbCl_3 , and AsCl_3 . II. The system $\text{SnCl}_4\text{-CCl}_3\text{COOH}$ and $\text{SnCl}_4\text{-CHCl}_2\text{COOH}$." by T. Shimizu and M. Umemoto. (p.984)

SO: Journal of General Chemistry (Zhurnal Obshchei Khimii) 1951, Volume 21, No. 6

Chemical Abst.
Vol. 48 No. 4
Feb. 25, 1954
General and Physical Chemistry

Binary systems formed by SnCl_4 , SbCl_3 , and AsCl_3 . I. The SnCl_4 - CH_3COOH system. M. Usanovich, T. Sumarokova, and V. Glushko (Kazakh Acad. Sci.). *J. Gen. Chem. U.S.S.R.* 21, 1078-8 (1951) (Engl. translation); *Zhur. Obshchei Khim.* 21, 981-4 (1951); cf. *C.A.* 44, 9853a; 46, 9401i, 9402b. From singular max. observed in viscosity and cond. isotherms, and shrinkage in sp. vol., evidence was obtained of chem. interaction in mixts. of SnCl_4 and CH_3COOH . Detns. at 50, 60, and 70° gave no definite indications of the compn. of any compds. formed. II. The SnCl_4 - CCl_3COOH and SnCl_4 - CHCl_2COOH systems. T. Sumarokova and M. Usanovich (Kazakh Acad. Sci.). *J. Gen. Chem. U.S.S.R.* 21, 1079-81 (1951) (Engl. translation); *Zhur. Obshchei Khim.* 21, 984-7 (1951). Viscosity and d. isotherms were detd. for the system SnCl_4 - CCl_3COOH at 50, 60, and 70°, and for the system SnCl_4 - CHCl_2COOH at 35, 50, 60, and 70°. Both systems are nonconductors. The absence of cond., the additivity of vols., and the smooth change of viscosity with compn. are taken as proof there is no chem. interaction in these systems. III. The SbCl_3 - CH_3COOH system. M. Usanovich and T. Sumarokova (Kazakh Acad. Sci.). *J. Gen. Chem. U.S.S.R.* 21, 1083-6 (1951) (Engl. translation); *Zhur. Obshchei Khim.* 21, 987-90 (1951). D., viscosity, and cond. values were detd. as functions of compn. in the SbCl_3 - AcOH system at 20, 50, and 60°. D. data show a slight vol. shrinkage in mixts. of the components. The viscosity isotherms exhibit max. at 52-3 mole % SbCl_3 , which shift toward higher SbCl_3 values at higher temps. A cond. max. at 70-80 mole % SbCl_3 shifts toward AcOH at higher temp. When cond. values are multiplied by the viscosity of the same solns., the cor. cond. isotherms all attain a max. at a compn. corresponding to $2\text{SbCl}_3 \cdot \text{AcOH}$. The temp. coeff. of cond. exhibits a max. at the compn. corresponding to $\text{SbCl}_3 \cdot \text{AcOH}$. The latter compd. was reported by B. N. Menshutkin (*C.A.* 6, 1280). The structures of the compds. found are given as $(\text{SbCl}_3 \cdot \text{AcOH})^+ \text{Cl}^-$ and $(\text{SbCl}_3 \cdot \text{AcOH})^+ \text{SbCl}_3^-$, the latter existing only in the liquid phase. Bernard M. Zeffert.

1155

Binary systems formed by SbCl_3 , SbCl_5 , and AsCl_3 . IV. The system $\text{SbCl}_3\text{-CH}_2\text{ClCOOH}$. M. I. Usanovich and T. Sumarokova. *Zhur. Obshch. Khim.* 21, 1214-15 (1951); cf. C.A. 45, 1787c. — The elec. cond., viscosity, and d. of the system $\text{SbCl}_3\text{-CH}_2\text{ClCOOH}$ were detd. at 50, 60, and 70° for changing values of the concn. of the components. Existence of the compds. $\text{SbCl}_3\text{CH}_2\text{ClCOOH}$ and $2\text{SbCl}_3\text{-CH}_2\text{ClCOOH}$ is indicated. The fusion diagram for the system shows the presence of a compd. of equimolar compn. V. The system $\text{SbCl}_3\text{-CCl}_3\text{COOH}$. T. Sumarokova and M. Usanovich. *Ibid.* 1219-22. — The elec. cond., viscosity, and d. of the system $\text{SbCl}_3\text{-CCl}_3\text{COOH}$ were detd. at 50, 60, and 70° for different concns. of the components. The elec. cond.-compn. and viscosity-compn. diagrams show that an acid-base reaction occurs between the components. The presence of the compds. $\text{SbCl}_3\text{CCl}_3\text{COOH}$ and $2\text{SbCl}_3\text{-CCl}_3\text{COOH}$ is also indicated. The fusion diagram exhibits a max. corresponding to the compd. $\text{SbCl}_3\text{CCl}_3\text{COOH}$ with a m.p. of 58°. J. Rovtar Leach

Lab. Phys. Chem., Inst. Chem. Sci., A S Kazakh SSR

SUMARKOVA, I.

"Binary systems containing SnCl_4 , SbCl_3 and AsCl_3 . V. The system $\text{SbCl}_3 - \text{CCl}_3\text{COOH}$."
T. Sumarkova and M. Usanovich. (p. 1219)

SO: Journal of General Chemistry (Zhurnal Obshchei Khimii) 1951, Vol 21, No 7.

CA

The binary systems constituted by stannic chloride, antimony trichloride, and arsenic trichloride. VI. The arsenic trichloride trichloroacetic acid system. T. Sumatohara and A. Rabikov. *J. Gen. Chem. U.S.S.R.* 21: 1531-4 (1951) (Engl. translation); *Zhur. Akad. Nauk* 21: 1375 (1951).—Measurement of the viscosity and d_4 at 25°, 35°, and 60° reveals no interaction in the system $AsCl_3$ - CCl_3COOH . The components are mutually sol. at all temps. at 60°, from 100 to 22.1 mole % $AsCl_3$ at 35°, and from 100

to 50.2 mole % at 25°. Limiting viscosities and d_4 are as follows:

Mole %	$\eta \times 10^4$	d_4	$\eta \times 10^4$	d_4	$\eta \times 10^4$	d_4
$AsCl_3$						
100	12.26	2.1624	10.50	2.1231	8.65	2.0824
39.21	33.44	1.8279	25.92	1.8043	13.93	1.7710
22.11	crystals	crystals	40.39	1.7256	22.76	1.6972
0	"	"	crystals	crystals	30.65	1.6951

1887

SHUMILOVA, T.

USSR/Chemistry - Arsenic Compounds

Aug 51

"Binary Systems Formed by SnCl_4 , SbCl_3 , and AsCl_3 .
VII. The Systems AsCl_3 - CH_2ClCOOH and AsCl_3 -
 CH_3COOH ," T. Sumarokova, T. Glushchenko, Student,
Lab of Phys Chem, Inst of Chem, Acad Sci Kazakh
SSR

"Zhur Obshch Khim" Vol XXI, No 8, pp 1376-1380

In system AsCl_3 - CH_2ClCOOH , form of isotherms of
viscosity and density at temps 50, 60, 70°C and
absence of elec cond showed that components do not
interact. In system AsCl_3 - CH_3COOH , study of elec
cond at temps 50, 60 and viscosity and density at
20, 50, 60, 70° established that there is acid-base
interaction.

1887

SUMAROKOVA, T.N.

SUMAROKOVA, T.N.; LITVYAK, I.O.

Complex compounds $\text{SnCl}_4 \cdot 2\text{A} \cdot \text{B}$ and $\text{SnCl}_4 \cdot 2\text{A} \cdot 2\text{B}$. Report no.1.
Izv.Sekt.plat.i blag.met. no.27:127-136 '52. (MLRA 7:5)

1. Institut khimicheskikh nauk Akademii nauk Kazakhskoy SSR,
Alma-Ata. (Compounds, Complex) (Tin compounds)

SUMAROKOVA, T. N.

SUMAROKOVA, T.N.; MAKSAI, L.I.

Complex compounds $\text{SnCl}_4 \cdot 2\text{A} \cdot \text{B}$ and $\text{SnCl}_4 \cdot 2\text{A} \cdot 2\text{B}$. Report no.2.
Izv. Sekt. plat. i blag. met. no. 27:137-151 '52. (MLRA 7:5)

1. Institut khimicheskikh nauk Akademii Kazakhskoy SSR, Alma-Ata.
(Compounds, Complex) (Tin compounds)

Chemical Abst.
Vol. 43
Apr. 10, 1954
Inorganic Chemistry

Binary systems formed by stannic chloride with fatty acids. I. *A. E. Yakovleva and T. Sumarokova. *Izv. Akad. Nauk Kazakh. S.S.R. No. III, 57, Kazan, No. 6, 30-31 (1953).*—Binary mixts. of SnCl_4 with EtCO_2H , PrCO_2H , and AmCO_2H were examd. by cond., d., and viscosity methods. The results, which are given graphically for 30°, 50°, and 70°, show the existence of complexes: $\text{SnCl}_4 \cdot 2\text{EtCO}_2\text{H}$, $\text{SnCl}_4 \cdot 3\text{EtCO}_2\text{H}$, $\text{SnCl}_4 \cdot 2\text{PrCO}_2\text{H}$, $\text{SnCl}_4 \cdot 3\text{PrCO}_2\text{H}$, $\text{SnCl}_4 \cdot 2\text{AmCO}_2\text{H}$, and $\text{SnCl}_4 \cdot 3\text{AmCO}_2\text{H}$. *Ref. 54-48.*—The work is extended to SnCl_4 systems with $\text{C}_{12}\text{H}_{25}\text{CO}_2\text{H}$ and $\text{C}_{18}\text{H}_{37}\text{CO}_2\text{H}$. Evidence is found for formation of complexes with 2 and 3 moles of the org. acids per mole of SnCl_4 . Previous work on complexes of substituted acids (cf. C. 48, 1787) is summarized. AcOH and chloro derivatives behave similarly in respect to SnCl_4 , SbCl_5 , AsCl_3 , HClO_4 , H_2SO_4 , and HNO_3 ; i.e., the action of protic and aprotic acids is similar. Concepts of acids and bases are discussed. G. M. Kovaleva

YAKOVLEVA, F.; SUMAROKOVA, T.

Binary systems formed by SnCl_4 with carboxylic acids of the aliphatic series.
Report no.2. Izv. AN Kazakh. SSR no.118:54-68 '53. (MLBA 6:10)
(Systems (Chemistry)) (Stannic chloride) (Carboxylic acids)

SUMAROKOVA, T.N.

4

✓ The existence of antimony tetrachloride. M. I. Usanovich, T. N. Sumarokova, and M. B. Beketov. *Izvest. Akad. Nauk Kazakh. S.S.R. No. 123, Ser. Khim. No. 7, 3-8(1953).*—Cond., viscosity, and d. of the system SbCl_3 - SbCl_5 were examd. at 50°, 60°, 70°, and 80°. The viscosity-compn. diagram shows smooth isotherms, whereas the sp. cond. compn. diagram shows a max. at about 15 mol. % SbCl_3 ; cor. sp. cond. declines with rise in temp. The results are best attributed to formation of the ion pair $(\text{SbCl}_4)^+ (\text{SbCl}_6)^-$. Cf. Weiland and Schmid, *Ber.* 38, 1059 (1905); 36, 541 (1903). G. M. Kosolapoff

SUMAROKOVA, T.; BOLIYALOV, K.

Oxonium compounds of esters with organic acids. Part 2.

System: cetyl acetate--acetic acid. Zhur.ob.khim. 25


no.3;477-479 Mr '55

(MLRA 8:6)

1. Institut khimicheskikh nauk Akademii nauk Kazakhskoy SSR
(Acetic acid)(Cetyl acetate)

"APPROVED FOR RELEASE: 08/26/2000

CIA-RDP86-00513R001653910020-4



APPROVED FOR RELEASE: 08/26/2000

CIA-RDP86-00513R001653910020-4"

"APPROVED FOR RELEASE: 08/26/2000

CIA-RDP86-00513R001653910020-4

Exhibit 1 - 1/1/1980

APPROVED FOR RELEASE: 08/26/2000


CIA-RDP86-00513R001653910020-4"

[illegible]

Instr Chem Sci AS tanggal 55R

"APPROVED FOR RELEASE: 08/26/2000

CIA-RDP86-00513R001653910020-4



APPROVED FOR RELEASE: 08/26/2000

CIA-RDP86-00513R001653910020-4"

SUMAROKOVA, T.; YARMUKHAMEDOVA, E.

Complex compounds of tin chloride and tin bromide with glycine.
Zhur.ob.khim. 26 no.12:3295-3301 D '56. (MLRA 10:7)

1. Institut khimicheskikh nauk Akademii nauk Kazakhskoy SSR.
(Tin chlorides) (Tin bromides) (Glycine)

S. SUMAROKOVA, T.N.

137-58-2-3907

Translation from: Referativnyy zhurnal, Metallurgiya, 1958, Nr 2, p 233 (USSR)

AUTHOR: Sumarokova, T.N.

TITLE: Eutectic Diagrams of the Fusibility of Binary Systems. Communication Nr 1 (Ob evtekticheskikh diagrammakh plavkosti dvoynykh sistem. Soobsheniye 1)

PERIODICAL: Izv. AN KazSSR. Ser. khim., 1957, Nr 1, pp 3-11

ABSTRACT: A check was made of the applicability of the Schroeder equation to real systems (organic substances, metals, salts) on the basis of available experimental data from the literature. It is shown that the Schroeder equation is valid at the most varied concentrations in different systems, and that many real systems are subject to the laws of ideal solutions. In ideal binary systems containing a common component, the liquidus curve is the geometric locus of the eutectic points. It is shown that molecular weight may be determined from the data of thermal analysis and the Schroeder equation.

A. F.

Card 1/1

1. Molecular weight—Determination 2. Schroeder equation—Applications

SUMAROKOVA, T.N.; PERMINOVA, D.

Eutectic fusibility diagrams of binary systems. Report no.2.
Izv. AN Kazakh. S.S.R. Ser. khim. no. 1:12-18 '57. (MLBA 10:5)
(Eutectics) (Systems (Chemistry))

SUMAROKOVA, T.; LITVYAK, I.

Complex tin compounds. Part 4. Zhur.ob.khim. 27 no.5:1125-1130
Zhur.ob.khim. 27 no.5:1125-1130 My '57. (MLBA 10:8)

1. Institut khimii Akademii nauk Kazakhskoy SSR.
(Complex compounds)
(Tin organic compounds)

AUTHORS: Sumarokova, T., Nevskaya, Yu.

79-12-41/43

TITLE: Complex Compounds of SnCl_4 , SnBr_4 and TiCl_4 With
Cineole (Kompleksnyye soedineniya SnCl_4 , SnBr_4 i
 TiCl_4 s Tsineolom).

PERIODICAL: Zhurnal Obshchey Khimii, 1957, Vol. 27, Nr 12,
pp. 3375-3379 (USSR)

ABSTRACT: The complex compounds SnCl_4 , SnBr_4 and TiCl_4 with organic
oxides have been little investigated. Compounds of SnCl_4 and
 SnBr_4 with dioxane, as well as of SnCl_4 with lactones
(SnX^2 2 A) are described in publications. The authors were
interested in the systematical investigation of the complex
compounds Sn^{+4} and Ti^{+4} with such organic compounds, which
possess in their molecule the group $-\text{COC}-$. They chose
cineole, which is a constituent part of many etheric oils.
The complex compounds of the tin- and titanium halides with
cineole were of an extended interest, in as much they are
connected with the research to find a method for the
quantitative determination of cineole. Mixtures of
 SnCl_4 , SnBr_4 and TiCl_4 in an exact molecular ratio with

Card 1/3

Complex Compounds of SnCl_4 , SnBr_4 and TiCl_4 With Cineole 79-12-41/43

cineole were prepared for the synthesis of these compounds. In the course of the reactions a considerable amount of heat was liberated. The components were mixed in indifferent solvents. The composition of the complex compounds was determined analytically and according to the cryoscopic method of titration. The amount of tin and titanium was computed as SnO_2 and TiO_2 the halides were determined according to Vollhard and the Cryoscopic measurements were conducted according to Beckmann. The authors put up diagrams on the basis of the results, from which the dependence of the depression (or of the molecular weight) on the composition, expressed in molecular percent, may be read. The following complex compounds were synthesized: $\text{SnCl}_4 \cdot 2\text{C}_{10}\text{H}_{18}\text{O}$, $\text{SnBr}_4 \cdot 2\text{C}_{10}\text{H}_{18}\text{O}$ and $\text{TiCl}_4 \cdot 2\text{C}_{10}\text{H}_{18}\text{O}$. The compound $\text{SnBr}_4 \cdot 2\text{C}_{10}\text{H}_{18}\text{O}$ possesses quite extraordinary properties: It shows a molecular weight of 746.8 and distillates already at 35°C . There are 3 figures, and 7 references, 5 of which are Slavic.

Card 2/3

Complex Compounds of SnCl_4 , SnBr_4 and TiCl_4 With Cineole 79-12-41/43

ASSOCIATION: Institute of Chemistry AS Kazakh SSR
(Institut khimii Akademii nauk Kazakhskoy SSR).

SUBMITTED: October 31, 1956

AVAILABLE: Library of Congress

1. Complex compounds - Synthesis

Card 3/3

On the Electrolytic Dissociation of Tin
and Antimony Complex Compounds.

~~SECRET~~
20-2-34/67

acid and ethyl acetate, where the organic compound contained the isotope C^{14} . In each case $SnCl_4$ was carried to the anode as well as to the cathode. This harmonizes with the equations 1 and 2. While this paper was being written, two essays were published (Miskidzh'van; Kuz'mina and Vol'nov) in which the authors suggest own schemes of the electrolytic dissociation of complex compounds. These schemes have in common that the formation of complex compounds is meant to represent an incorporating reaction, i.e. the molecule of the organic matter is to be a component of the complex cation. The experiments of the authors of the present paper contradict these schemes. For only the complex compounds of the 3-halides of antimony (and arsenic) with organic oxygen containing substance are incorporation products. The compounds of the same organic substances with the 4-halides of tin are not incorporation products. In the case of electrolysis according to the equation 3 the authors of the present paper have added methyl-red in addition to the acid marked with C^{14} . As expected, the pigment moved into the same direction as the acetic acid, that is towards the cathode.
(With 8 citations from publications).

Card 2/3

POLOSUKHIN, Porfiriy Porfir'yevich, zasluzhenny master sporta. Prinsipal
uchastiye: REVZIN, Sergey Vladimirovich, inzh.-vozdukhoplavatel'.
SUMAROKOVA, T.N., red.; MANINA, M.P., tekhn.red.

[Notes of an amateur navigator and parachutist; as told to Sergei
Revzin] Zapiski sportsmena-vozdukhoplavatelya i parashyutista.
Literaturnaya zapis' Sergeia Revzina. Izd.3., dop. i perer.
Moskva, Gos.izd-vo "Fizkul'tura i sport," 1958. 230 p.

(MIRA 12:12)

(Polosukhin, Porfirii Porfir'evich, 1910-)

SUMAROKOVA T N

SUMAROKOVA T N

5(4)
APR 1950
TIT 151
PRAK 101010101
ABSTRACT

Yevl'yan, V. P., Korablina, V. B., 1947/53-54-5-30/50
Tsitimirovskiy, E. B.

Conference Discussion on the Methods of Investigating the
Complex Formation in Solutions (Sovetskaya-khimiya
po metodam khimicheskoy kompleksirovaniya v rastvorakh)

Investiya vreshnaya sobremennosti. Khimicheskaya
khimicheskaya khimicheskaya, 1946, No. 3, pp. 113-114 (USSR)
From February 10 to 21, 1949 a conference discussion took
place at the Institute of Chemistry, it dealt with the subjects
mentioned in the title. It was called in a decision of the
USSR All-Union Conference on the Chemistry of Complex
Formations. More than 200 persons attended the conference,
among them 103 delegates from various towns of the USSR.
At the conference methods of determining the composition of
the complexes in solutions were discussed, as well as the
methods of calculating the instability constants according
to experimental data and problems concerning the influence
of the solvent upon the processes of complex formation.

E. B. Yevl'yan, "Physical and Chemical Analysis of the
System With 5 Colored Complexes in the Solution", the results
of a systematic investigation in copper-pyrimidine-sulfate,
as well as in copper-pyrimidine-sulfate by means of the
optical method were dealt with. The results of the work of
Flicker, the idea of a further investigation of the complex
formation processes in solutions was developed. Besides the
determination of the composition of the complexes, the
also the physical and chemical properties, the chemical nature
and the structure of the complex compounds must be investi-
gated.

Card 2/16

1. A. I. Kikhezer and E. B. Yevl'yan in their lecture "Investiga-
tion of the Polymerization of Iso-Poly Acids in Solutions"
mentioned experimental results of the investigation of the
polymerization in solutions of polyacetic acid. The authors
proved that especially the polyacetic acid within a certain
range of the pH values and the concentrations exists as a
number of compounds that can be expressed by an overall formula
 $HO_2C(CH_2COO)_nH$. In the lecture by E. V. Kiselev and V. J.

Card 4/16

Spiridonov investigation results on basic salts taking into
account the complex formation in solutions by means of the
potentiometric method were mentioned. The authors with aim,
sodium and iodine. In the evaluation of their results the
authors employed the method of the basic difference. The
calculation of the consecutive constants was carried out
according to the interpolation formula by Benet, E. A.
Chapalinsky held a lecture on "pH Measurement Method of the
Solutions in Combination With the System $Cu^{2+}-HCl - H_2O$ in Investi-
gating Complex Copper Compounds in Saturated Solutions". It
was found that the substance at the bottom of the liquid is
more basic than the solution. Furthermore, the increased
acidity of the solution from the viewpoint of the formation
of hydroxy-chloro complexes in the solution was explained.
V. I. Kiselev opened the discussion with his lecture. He
pointed out the necessity of utilizing the concepts sorted
out in the investigations of the polymerization in organic
chemistry in the chemistry of polynuclear complexes. A. A.
Grisberg thinks that the new approach of the hydrolysis

Card 5/16

Conference Discussion on the Methods of
Investigating the Complex Formation in Solutions

507/153-54-3-10/30

Investigation as developed by the Scandinavian school is of high value. He also pointed to the necessity of studying the kinetics of the polymerization process and a quantitative determination of the strength of the polymers. A. E. Babko pointed out that the study of the polymer structure was necessary. E. P. Kozlov mentioned in his lecture that the rather widely spread polymerization type according to the mechanism of chain growth is not obtained in all cases. The following scientists took part in the discussion: V. E. Zaslavskiy, V. E. Zaslavskiy, I. V. Tsvetkov, and E. A. Yatskovskiy. A. E. Babko took part in his lecture "Methods of Determining the Dissociation Constants of the Complex Groups in Solutions". E. P. Kozlov discussed in his lecture "Calculation Methods of the Instability Constants of the Complex Compounds According to Experimental Data" the possibilities of using the known calculation methods of the instability constants for various cases of the complex formation in solution. If several monomolecular complexes are formed the displacement method by Abegg and Bedender (completed by A. E. Babko) cannot be recommended for the calculation of the instability constant. The lecturer discussed the displacement method of the polyamides proposed by B. Jernum, Beden, Kozlov, Matshard, Edolloy and other authors. The constants calculated in this way are not very accurate. It was proved that the method of successive approximations can be used in the case of the complex formation. The most probable value of the physical constants can be obtained by the method of the least squares. E. P. Kozlov, V. E. Zaslavskiy and E. A. Yatskovskiy described the determination of the instability constants of the complex compounds of the uranum and iron which are based on the investigation of the equilibrium displacement of the complex formation by silver ions. E. E. Bol'shakov, I. V. Tsvetkov and G. S. Kozlovskiy held a lecture on "The Role of the Time Factor in the Investigation of the Complex Formation". In the discussion on the lecture A. A. Grinberg mentioned that due to the slow adjustment of the equilibria the methods discussed of determining the instability constants (palladium and cobalt complexes) can often not be employed. A. V. Ablov pointed out the necessity of deriving direct methods of proving the existence of intermediate forms in a step-wise complex formation. E. E. Bol'shakov mentioned that the instability constants of slowly dissociating complexes can be calculated from thermodynamic data. L. P. Adamevich, A. M. Golub and others took part in the discussion on the lecture. A. E. Babko requested inclusion in the next conference on the chemistry of complex compounds a lecture in which various calculated by methods of the instability constants should be compared with the results of experimental data can lead. E. P. Kozlov stressed that in the determination of the instability constants all chemical equilibria should be taken into account that render complex the complex formation process in the solution, especially the hydrolysis processes of the central ion and the addendum. In the lecture delivered by V. E. Zaslavskiy and A. P. Zaslavskiy "Application of the Distribution Method to the Investigation of the Stability Constants

Card 6/16

Card 7/16

Card 8/16

Conference Discussion on the Methods of
Investigating the Complex Formation in Solutions

20/153-54-3-30/20

of Some Thorium Complex Compounds" results obtained from the experimental investigation of the distribution of thorium compounds in the systems: acetylacetone - benzene - water, and 2-oxo-1,4-naphthoquinone - chloroform - water were given. From these data the instability constants of the thorium complexes with acetylacetone and 2-oxo-1,4-naphthoquinone were calculated. I. V. Tsvetkov, G. B. Zhuravskiy and V. I. Gerasimov held a lecture on the application of the solubility method in the determination of the stability of complex compounds in solutions. In this lecture the other methods of investigating complex formation phenomena in the solution were discussed (pH method, potentiometric method, etc.). I. V. Tsvetkov held a lecture on the "Application of the Solubility Method in Studying the Pathological Complexes of Metals". He used the determined quantitative characteristics of the reaction of the transition of the pathological complexes of cobalt, nickel, copper and zinc, as well as of the free pathological complex into the sulfuric acid solution for the theoretical reasoning, and as an experimental proof of the existence of complexes in the complexes investigated. These characteristics also served him as a proof of new electrochemical formulas of pathological complexes and its complex derivatives. In the lecture delivered by I. L. Krupatkin on "The Method of the Free Solvent as a Method of Investigating the Formation and Properties of Organic Complexes" it was proved that this method makes it possible to determine the number of complexes formed in the system, their composition and relative stability. V. I. Zhuravskiy, A. K. Babko, E. P. Kozlov, I. S. Buzhik and I. A. Tsvetkov took part in this discussion. In the lecture delivered by A. A. Grishin and E. P. Kozlov on the complex formation in the systems of the coordination compounds of palladium compounds (II) with a coordination number 4 and 6, it was proved that in the case of the complexes of palladium in these systems the coordination number 4 and 6 are estimated. I. P. Kozlovish mentioned a new manipulation in the spectrochemical investigation of the complex compounds that can be used in systems with the formation (or predominance) of one single complex. This method makes it possible to determine the composition and instability constant of the complex. In the lecture delivered by E. B. Yatsinskaya and V. I. Gerasimov on the application of the theory of crystal field to the determination of the composition and structure of the chloride complexes of cobalt, nickel and copper, according to the absorption spectra of these complexes was discussed. It was proved that in a hydrochloric acid solution above 5 mole/liter in the solution there exists an equilibrium between the tetrahedral and octahedral forms of the cobalt chloride complexes. Yu. P. Kozlovish proved in his lecture "The Application of Radioactive Isotopes in the Investigation of the Solvation Equilibrium in Solutions of Complex Compounds" the possibility of using data on the isotope exchange to clarify the structure of the complexes and mechanism of the hydration processes. I. Kozlovish mentioned in his lecture the use of radioactive isotopes in the study of tin and antimony complexes. In the lecture delivered by A. V. Ahlov, V. I. Tsvetkov, E. P. Kozlov and A. M. Golub took part in the discussion of the lecture. The usefulness of applying the theory of the crystalline fields in explaining the results obtained from the absorption spectra of the com-

Card 9/16

Card 10/16

Card 11/16

Conference Discussion on the Methods of Investigating the Complex Formation in Solutions

06-05-1935-3-30/20

plex compounds was stressed. In the lecture delivered by I. A. Babko on "The Investigation of the Complex Formation of the Solvent on the Basis of the Dielectric Permeability and the Polarization" the principal methods mentioned were presented. Then the principal method for investigating the compounds of this nature, the "diffusion" products. The lecture delivered by the type book and Ye. Ye. Kriss "Employing the Method of the Dielectric Constant for Investigating Complex Compounds of the Type of Crystalline Solvents in Solutions" dealt with the investigation of the solvates of lanthanum and cerium chlorides with benzene, as well as with the study of the compounds formed in heterogeneous systems with tributyl phosphate and acetic acid. V. V. Korotkov gave in his lecture "The Polarographic Method of Investigating the Complex Formation in Solutions" a survey of the applications of the polarographic method in the study of the complex compounds, and illustrated several fine characteristic features of this method. The lecture delivered by T. S. Shadrin on "The Cryoscopic Method of Investigating the Complex Formation in Solutions" surveyed of the possibilities of the cryoscopic method and its application in the study of several complex compounds of the type of crystalline solvents. The lecture delivered by A. K. Golub described the results of his investigations of thioetheric compounds of several types. A. A. Plakhot and Ya. Ye. Pliashov on the lecture "The Cryoscopic Method of Investigating Thioetheric Compounds" to be of considerable value. K. I. Yermolov pointed out that the publication of the surveys on individual methods of investigating the complex formation and the results of the investigations of the complex formation reactions would be desired. This concerns especially the polarographic method. The cryoscopic method should be brought to a level that makes the calculation of the equilibrium constants of the processes to be investigated possible. The problem of the method of evaluating the experimental results becomes more and more important. Many scientists use the insolubility constants without taking into account the way in which they had been obtained. The calculation methods employed by A. K. Golub are an step back, as compared to those employed at present. In his lecture K. I. Yermolov pointed out the extremely great importance of the mathematical evaluation of the results obtained, as well as of the plotting of curves. A. K. Babko suggested selecting one or two systems that are experimentally well investigated, and to evaluate the results obtained according to different methods so that it is possible to check and evaluate them. Ye. I. Tur'yan took part in the discussion. Ya. A. Fikhter discussed in his lecture "The Effect of the Solvent on the Complex Formation Processes as Well as on the State of Equilibrium in the Solutions of Complex Compounds" the influence exerted by the solvent upon the molecular state, upon the solvation of the system components, upon the stabilization of the complexes formed by the system, upon the stepwise dissociation of the complexes and upon the change in the dielectric constant of the solution. By the dielectric constant upon the complex formation process was discussed. It was concluded that a direct relation does not exist, and that the chemical nature of the solvent must be taken into account. A. V. Akher and L. V. Segurenko held a lecture on "The Spectroscopic Investigation of Nickel Carbonyl in Various Solvents". The invariability constants of the complexes were determined and it was proved that the

Card 12/16

Card 13/16

9/1/11 PM

Conference Discussion on the Methods of
Investigating the Complex Formation in Solutions

30/1/55-56-5-30/50

stability of the 'pyridinate' is changed in dependence on the solvent. In I. I. Tsv'yan's lecture "The Influence of the Solvent Upon the Competition and Stability of Complexes" discussed the polarographic investigation method of the chlorides and thiocyanate complexes of lead in aqueous ethanol solutions at different content of the non-aqueous solvent and at a constant ionic strength. A step-wise character of the complex formation was found as well as the instability constants of the complexes. The influence of the dielectric constant of the solution on the stability of the investigated complexes was studied. In the lecture by V. I. Kuznetsov, "Investigation of Aqueous Complexes in Aqueous Solution" the main attention was devoted to the accuracy of the qualitative recording of the solvation effects in the complex formation. The applicability of the polarographic method in the determination of the competition and stability of the aqueous complexes in mixed solvents was proved and experimental material on the thermodynamics of the dissociation of the cadmium-aqueous complexes in aqueous ethanol solutions was mentioned. V. E. Tel'macher, V. I. Kuznetsov

Card 15/16

and I. V. Tsanayev stressed in their lectures the necessity of a more complete and general investigation of the solvation processes. A. E. Babko and A. M. Golub pointed out the great importance of the investigations of the complex formation equilibria in non-aqueous solutions, and made several critical comments on the lecture by I. I. Tsv'yan. The following scientists took part in this discussion: L. P. Demerits, O. I. Khokhlovskiy, A. F. Koshits and A. G. Shatshber. At the final meeting of the conference A. A. Krimberg, corresponding member, AS USSR, said in his speech that the conference discussed the solvation effects in the determination of the competition and stability of the complexes, as well as of the method used in the study of the quantitative characteristics of the stepwise complex formation was extremely useful for all who attended this conference.

Card 16/16
CONF-55-50976

AUTHORS: Molodtsov, T., Smirnov, T. SOV/ 78-3-7-33/44

TITLE: The Systems $PbCl_2-PbBr_2$ and $PbBr_2-PbI_2$ (Sistemy $PbCl_2-PbBr_2$ i $PbBr_2-PbI_2$)

PERIODICAL: Zhurnal neorganicheskoy khimii, 1958, Vol. 3, Nr 7, pp 1655-1660 (USSR)

ABSTRACT: The systems $PbCl_2-PbBr_2$ and $PbBr_2-PbI_2$ were investigated by means of thermal analysis. The compound $PbBr_2$ was found to occur in form of two modifications. At $344^\circ C$ a phase change $\alpha \rightleftharpoons \beta$ occurs. The melting temperatures of the β -modifications of $PbCl_2 = 480^\circ C$, $PbBr_2 = 364^\circ C$ and $PbI_2 = 396^\circ$ were determined. The solid solutions of $PbCl_2$ and $PbBr_2$ indicate the existence of the equimolar compound $PbClBr$. The phase transformation of liquid solutions into solid α -modification and the transformation of solid solutions of the α -modification into solid solutions of the β -modification characterize the melting diagram of the systems. In the system $PbBr_2-PbI_2$ there also exist α - and β phase changes. There are 2 figures, 2 tables and 11 references, 6 of which are Soviet.

Card 1/2

The Systems $PbCl_2-PbBr_2$ and $PbBr_2-PbI_2$

SOV/78-3-7-33/44

ASSOCIATION: Imambetovskiy AN Kazakhskoy SSR (Institute of Chemistry, AS
Kazakhskaya SSR)

SUBMITTED: June 17, 1977

1. Lead bromide-lead iodide systems--Analysis 2. Lead bromide
-lead chloride systems--Analysis

Card 2/2

AUTHORS: Yarmukhamedova, E. Sh., Sumarokova, T. N. 79-28-5-67/69

TITLE: Complex Compounds of Tin Chloride and Tin Bromide With Urea
(Kompleksnyye soedineniya khlornogo i bromnogo olova s mochevinoy)

PERIODICAL: Zhurnal Obshchey Khimii, 1958, Vol 28, Nr 5,
pp 1410 - 1412 (USSR)

ABSTRACT: In the systematic investigation of complex compounds of halides of tetravalent tin with organic compounds containing nitrogen and oxygen, the authors found that thiourea enters reaction with tin chloride and tin bromide with formation of a complex compound of the composition $\text{SnX}_{4.2}(\text{NH}_2)_2\text{CS}$ (Reference 1). It was of interest for the authors to experience by investigations which way urea would react on the halides of tetravalent tin. Taking into account the similarity of urea with thiourea it was assumed that also the former had to form compounds of the same kind with tin chloride and tin bromide. Tin chloride was purified by repeated distillation and the fraction with the boiling temperature 109°C (690°mm) was stored in sealed ampoules. Tin bromide was treated the same way. The complex com-

Card 1/2

79-28-5-67/69

Complex Compounds of Tin Chloride and Tin Bromide With Urea

pound of tin chloride with urea was obtained by direct action of tin chloride on it without solvent - that of tin bromide the same way. Thus the complex compounds of tin chloride and tin bromide with urea of the following composition were synthesized: $\text{SnCl}_4 \cdot 2(\text{NH}_2)_2\text{CO}$ and $\text{SnBr}_2 \cdot 2(\text{NH}_2)_2\text{CO}$. These complex compounds are crystalline products, do not change in air or dissolve easily in organic media. There are 2 figures and 1 reference, 1 of which is Soviet.

ASSOCIATION: Institut khimicheskikh nauk Akademii nauk Kazakhskoy SSR
(Institute for Chemical Sciences, AS Kazakh SSR)

SUBMITTED: April 19, 1957

Card 2/2

AUTHORS: Sumarokova, T. N., Arsenov, G. I.

76-32-5-32/47

TITLE: Methods of Cryoscopic Measurements (K metodike krioskopicheskikh izmereniy)

PERIODICAL: Zhurnal fizicheskoy khimii, 1958: Vol. 32, Nr 5, pp.1153-1154 (USSR)

ABSTRACT: The usual method of determination based on the measurement of the maximum temperature rise gives wrong results, as can be seen from a given diagram, as the measuring values do not correspond to the real freezing point of the solutions. In determinations according to M. Bakeyev (Ref 2) which are based on the determination of the heating curves the error by undercooling is removed, however, errors due to insufficient mixing occur. An electromagnetic arrangement of mixing is described, using a multivibrator with two lamps L₁-6Zh8 and L₂-6P3 as well as a transformer ELS-2. In order to improve the distribution of the crystals the cryoscope has an elevated bottom. In order to determine the freezing temperature the heating curves are taken by plotting the temperature on the ordinate and the time on the abscissa. By

Card 1/2

76-32-5-32/47

Methods of Cryoscopic Measurements

means of an example, a solution of benzylalcohol in benzene, the obtained diagrams are represented, the melting temperatures (crystallization) being determined by the point of intersection of two straights. The described electromagnetic mixer can also be used for other purposes. There are 4 figures and 3 references, 3 of which are Soviet.

ASSOCIATION: Akademiya nauk Kazakhskoy SSR, Institut khimii, Insitut energietiki
(Institutes of Chemistry and Power Engineering, AS Kazakhskaya SSR)

SUBMITTED: May 3, 1957
1. Liquids--Freezing 2. Freezing points--Determination

Card 2/2

GUKOV, Valentin Ivanovich; SUMAROKOVA, T.N., red.; FEKLISOVA, T.D.,
tekhn.red.

[In the land of untouched treasures] V kraiu netremutnykh
sokrovishch. Moskva, Gos.izd-vo "Fizkul'tura i sport,"
1959. 86 p. (MIRA 12:6)
(Altai Territory---Description and travel)

SOV/79-29-5-5/75

5(4)
 AUTHORS: Sumarokova, T., Omarova, R.
 TITLE: On the Interaction of Tin Chloride With Esters. 1. (O vzaimodeyst-
 vii khlornogo olova so slozhnymi efirami.1.)
 PERIODICAL: Zhurnal obshchey khimii, 1959, Vol 29, Nr 5,
 pp 1430 - 1437 (USSR)
 ABSTRACT: In the present paper the systems $\text{SnCl}_4 - \text{CCl}_3\text{COOC}_2\text{H}_5$ and
 $\text{SnCl}_4 - \text{CH}_2\text{ClCOOC}_2\text{H}_5$ by means of physico-chemical analysis
 conductionometry, viscosimetry, volumetry and cryoscopy were inves-
 tigated. The operational methods, the preparation and purifica-
 tion of tin chloride were described previously (Refs 12 and 16).
 The results obtained by determination of the viscosity and the
 values B calculated from the equation

$$\eta = A e^{\frac{B}{RT}}$$
 (Refs 17-19) are presented in table 1. Table 2 gives
 the determination results of the density. In figure 1 the diagrams
 property - composition of the system $\text{SnCl}_4 - \text{CCl}_3\text{COOC}_2\text{H}_5$ are
 shown. The determination results of viscosity, specific conducti-

Card 1/3

On the Interaction of Tin Chloride With Esters.1.

SOV/79-29-5-5/75

vity and density for the system $\text{SnCl}_4 - \text{CH}_2\text{ClCOOC}_2\text{H}_5$ are given in table 3 and 4, the calculated values of the corrected conductivity, temperature coefficient of the conductivity, and of the constant B are presented in table 5. Figure 2 gives the diagrams property - composition of this system. On investigating the behavior of tin chloride with respect to ethyl trichloro- and ethyl monochloro-acetate the following was found: In the system $\text{SnCl}_4 - \text{CCl}_3\text{COOC}_2\text{H}_5$ the components react with one another until the compound $\text{SnCl}_4 \cdot 2\text{CCl}_3\text{COOC}_2\text{H}_5$ is formed. In this system the electric conductivity is practically not existing. In the system $\text{SnCl}_4 - \text{CH}_2\text{ClCOOC}_2\text{H}_5$ the reaction between the components is carried on up to the formation of the compounds $\text{SnCl}_4 \cdot 2\text{CH}_2\text{CClCOOC}_2\text{H}_5$ and $\text{SnCl}_4 \cdot 3\text{CH}_2\text{ClCOOC}_2\text{H}_5$. The latter determines the electric conductivity of the system. As can be seen from figures 4 and 5 the complex acids $\text{SnCl}_4 \cdot 2\text{RCOOH}$ are considerably stronger than $\text{SnCl}_4 \cdot 2\text{RCOOR}'$. From the comparison of

Card 2/3

On the Interaction of Tin Chloride With Esters. 1.

SOV/79-29-5-5/75

the behavior of ethyl- and ethyl-monochloro-acetate with the behavior of ethyl-trichloro-acetate with respect to tin chloride it can be seen that the electric conductivity then appears in the systems formed from tin chloride and esters, if the compound $\text{SnCl}_4 \cdot 3\text{RCOOR}'$ is formed in the solution. There are 5 figures, 5 tables, and 24 references, 19 of which are Soviet.

ASSOCIATION: Institut khimicheskikh nauk Akademii nauk Kazakhskoy SSR
(Institute of Chemical Sciences of the Academy of Sciences,
Kazakhskaya SSR)

SUBMITTED: April 18, 1958

Card 3/3

SOV/79-29-5-6/75

5(4)

AUTHORS: Sumarokova, T., Omarova, R., Kuz'menko, N.

TITLE: On the Interaction of Tin Chloride With Esters. 2. (O vzaimodeystvii khlornogo olova so slozhnymi efirami. 2.)

PERIODICAL: Zhurnal obshchey khimii, 1959, Vol 29, Nr 5, pp 1437 - 1442 (USSR)

ABSTRACT: In the present paper the influence exercised by the length of the aliphatic alcohol radical upon the acid-basic properties of esters was investigated. Viscosity, density and electric conductivity of the systems $\text{SnCl}_4 - \text{CH}_3\text{COOC}_8\text{H}_{17}$ and $\text{SnCl}_4 - \text{CH}_3\text{COOC}_{16}\text{H}_{33}$ were investigated. The results obtained on the determination of the properties as well as the calculated temperature coefficients of the electric conductivity, the corrected conductivity and the constant B for the system $\text{SnCl}_4 - \text{CH}_3\text{COOC}_8\text{H}_{17}$ which was investigated at 25 and 50°, are given in table 1. In figure 1 the diagrams ~~property~~ - composition are presented. By physico-chemical analysis it could be concluded that the components of the system react with one another, thus forming a complex compound $\text{SnCl}_4 \cdot 2\text{CH}_3\text{COOC}_8\text{H}_{17}$. The system $\text{SnCl}_4 - \text{CH}_3\text{COOC}_{16}\text{H}_{33}$ was in-

Card 1/3

On the Interaction of Tin Chloride With Esters. 2.

SOV/79-29-5-6/75

vestigated at 40, 50, 60 and 70°. The determination results are listed in tables 2 and 3, the calculated values of the corrected electric conductivity, the temperature coefficient of the conductivity and the constant B in table 4. The diagrams property-composition can be seen in figure 2. On the strength of the physico-chemical analysis it could be concluded that a complex compound $\text{SnCl}_4 \cdot 2\text{CH}_3\text{COOC}_{16}\text{H}_{33}$ is formed in the system. It was separated in free state. Its melting point is 56°. The electric conductivity in the system indicates the existence of this compound. Figure 3 compares the values of the electric conductivity at 50° in the systems $\text{SnCl}_4 - \text{CH}_3\text{COOC}_2\text{H}_5$ (Ref 4), $\text{SnCl}_4 - \text{CH}_3\text{COOC}_8\text{H}_{17}$ and $\text{SnCl}_4 - \text{CH}_3\text{COOC}_{16}\text{H}_{33}$. The value of the corrected conductivity of cetyl acetate solutions is seen to be far smaller than in octyl acetate and especially ethyl acetate solutions. This can be explained by the fact that the formation of the complex compounds $\text{SnCl}_4 \cdot 3\text{RCOOR}'$ is a secondary redox reaction. It proceeds via the stage of the formation of complex acids $\text{SnCl}_4 \cdot 2\text{RCOOR}'$ which become weaker on lengthening

Card 2/3

On the Interaction of Tin Chloride With Esters. 2.

SOV/79-29-5-6/75

of the radical. A similar rule was found in the systems formed from tin chloride and carboxylic acids (Refs 16,17). There are 3 figures, 4 tables, and 17 references, 13 of which are Soviet.

ASSOCIATION: Institut khimicheskikh nauk Akademii nauk Kazakhskoy SSR
(Institute of Chemical Sciences of the Academy of Sciences,
Kazakhskaya SSR)

SUBMITTED: April 18, 1958

Card 3/3

5(4)

AUTHOR:

Smirnov, T. E., Fialkov, Ya. A.,
(Alma-Ata), Deceased

SOV/76-33-1-31/45

TITLE:

On the Cryoscopic Method of the Physico-Chemical Analysis
(Classification of Diagrams) (O krioskopicheskom metode
fiziko-khimicheskogo analiza (klassifikatsiya diagramm))

PERIODICAL:

Zhurnal fizicheskoy khimii, 1959, Vol 33, Nr 1, pp 184-191
(USSR)

ABSTRACT:

The first method of this type has been worked out by
V. V. Udovenko (Refs 1-7) and is based on the establishment of
the function between the molecular weight of the components
(and their mixtures) of the concentration in the cryoscopic
solvent and the marking of the intersection points of the
isoconcentration in the coordinates molecular weight-
composition. For plotting the diagram composition-properties
by the cryoscopic method of the physico-chemical analysis of
the system A-B-solvent, N. A. Izmaylov (Refs 8-11) (and others
(Refs 12, 13)) suggested the establishment of the deviation
from the additive depression on the ordinate of properties.
Ya. A. Fialkov and I. D. Muzyka (Ref 16) used the depression
change as a property and based the determination on a

Card 1/3

On the Cryoscopic Method of the
Physico-Chemical Analysis (Classification of Diagrams)

SOV/76-33-1-31/45

measurement of the depression of the solutions which corresponds to the intersection point of a pencil of lines in the diagram depression change-composition (Fig 1). The use of precipitation reactions in cryoscopic investigations brought about a considerable increase in analytic possibilities (Ref 17). The amount of the depression is connected to the concentration by the equation according to Schröder (Shreder) (1). On account of (1) the diagram of the ideal case is established and the diagram types depression-composition of systems with a chemical reaction are investigated. It is stated that (in this case) there are three basic types of diagrams. The first type shows a decreasing depression value, the second type a constant value, and the third type a value increasing up to the end point of the depression. A number of examples is given illustrating the types mentioned

Card 2/3

SOV/76-55-1-51/45

On the Cryoscopic Method of the
Physico-Chemical Analysis (Classification of Diagrams)

(e.g. $\text{SnCl}_4\text{-C}_6\text{H}_4\text{O}_2$ in n-dichlorobenzene, $\text{TiCl}_4\text{-C}_4\text{H}_8\text{O}_2$
in benzene (Ref 19), $\text{NH}(\text{C}_2\text{H}_5)_2\text{-C}_3\text{H}_5\text{NCS}$ in benzene (Ref 16),
 $\text{NH}_2\text{CH}_2\text{COOH - CCl}_3\text{COOH}$ in acetic acid, etc). Several
deficiencies of a previous paper (Ref 8) are pointed out.
There are 3 figures and 21 Soviet references.

ASSOCIATION: Akademiya nauk KazSSR, Institut khimicheskikh nauk. Akademiya
nauk USSR Institut obshchey i neorganicheskoy khimii
(Academy of Sciences, Kazakhskaya SSR, Institute of Chemical Sciences
Academy of Sciences USSR Institute of General and Inorganic
Chemistry)

SUBMITTED: July 8, 1957

Card 3/3

SUMAROKOVA, T.M.

Shape of the liquidus curves of phase diagrams for systems
with chemical interaction. Zhur.neorg.khim. 5 no.7:
1572-1576 J1 '60. (MIRA 13:7)

1. Institut khimii Akademii nauk KazSSR.
(Systems(Chemistry))

SUMAROKOVA, T.N.; MODESTOVA, T.P.

Fusibility in the system $PbCl_2$ - KCl. Zhur. neorg. khim. 5 no.11:
2477-2482 N '60. (MIRA 13:11)

1. Institut khimii Akademii nauk Kazakhskoy SSR.
(Lead chloride) (Potassium chloride)